

tropenia associated with anorexia nervosa, marrow reserves are adequate and there is no increased incidence of associated opportunistic infections.¹⁷

Another possible cause of pulmonary injury in malnourished patients involves an associated antioxidant deficiency. Glutathione and other similar proteins are depleted in malnourished patients. In animal studies, increased susceptibility to oxygen toxicity occurs with starvation, especially at the oxygen concentrations required to oxygenate our patient.^{18,19}

Several behaviors associated with anorexia nervosa and bulimia predispose patients to pulmonary complications. Pneumomediastinum has been reported in association with bulimia.²⁰ Aspiration of foreign material is also a major concern. Though mentioned in reviews, there are no reports documenting this complication.⁴ Self-mutilative behavior without suicidal intent, in addition to alcohol and drug abuse, is frequently seen in bulimic patients.²¹ The circumstances surrounding our patient's lipid pneumonitis suggest a volitional component. The use of mineral oil as a laxative with accidental inhalation or aspiration is also possible.

Exogenous lipid pneumonia is an unusual pulmonary problem and has not been reported previously in association with anorexia nervosa.²² Though most often found as a slowly progressive, minimally symptomatic disease process, it occasionally presents as an acute pneumonitis.^{23,24} There is a high incidence of secondary infection due to plugging and impaired mucociliary clearance.^{25,26} Occasionally the pneumonitis progresses to interstitial fibrosis.²⁷ Respiratory failure is rare. Several factors mentioned previously may have contributed to this patient's pulmonary disease, although pathologically no other cause could be found. The use of corticosteroids has been suggested as a potential treatment.²⁸ With corticosteroid treatment, our patient has shown moderate improvement in lung function, suggesting resolution of the acute pneumonitis with residual fibrosis.

Pulmonary disorders in patients with anorexia nervosa are uncommon, though multiple complications can occur. With the frequent abuse of laxatives, including mineral oil, by anorexic patients, lipid pneumonia may develop more commonly than previously appreciated. The insidious nature of lipid pneumonia makes this an easily overlooked and potentially severe complication, which requires a high index of suspicion for prompt diagnosis.

REFERENCES

- Herzog DB, Copeland PM: Eating disorders. *N Engl J Med* 1985; 313:295-303
- Schwabe AD, Lippe BM, Chang RJ, et al: Anorexia nervosa. *Ann Intern Med* 1981; 94:371-381
- Harris RT: Bulimarexia and related eating disorders with medical complications. *Ann Intern Med* 1983; 99:800-807
- Brotman AW, Rigotti NA, Herzog DB: Medical complications of eating disorders. *Compr Psychiatry* 1985; 26:258-272
- Heysfield SB, Bethel RA, Ansley JD, et al: Cardiac abnormalities in cachectic patients before and during nutritional repletion. *Am Heart J* 1978; 95:584-594
- Gottdiener JS, Gross HA, Henry WL, et al: Effects of self-induced starvation on cardiac size and function in anorexia nervosa. *Circulation* 1978; 58:425-433
- Brotman AW, Stern TA: Case report of cardiovascular abnormalities in anorexia nervosa. *Am J Psychiatry* 1983; 140:1227-1228
- Hurd HP, Palumbo PJ, Gharib H: Hypothalamic-endocrine dysfunction in anorexia nervosa. *Mayo Clin Proc* 1977; 52:711-716
- Fohlin L: Body composition, cardiovascular and renal function in adolescent patients with anorexia nervosa. *Acta Paediatr Scand [Suppl]* 1977; 268:1-20
- Adler AG, Wolinsky P, Krall RA, et al: Death resulting from ipecac syrup poisoning. *JAMA* 1980; 243:1927-1928
- Spark RF, Arky RA, Boulter PR, et al: Renin, aldosterone and glucagon in the natriuresis of fasting. *N Engl J Med* 1975; 292:1335-1340
- Low DK, Dudrick SJ, Abdou NI: Immunocompetence of patients with protein-calorie malnutrition. *Ann Intern Med* 1973; 79:545-550

- Gotch FM, Spry CJF, Mowat AG, et al: Reversible granulocyte killing defect in anorexia nervosa. *Clin Exp Immunol* 1975; 21:244-249
- Kim Y, Michael AF: Hypocomplementemia in anorexia nervosa. *J Pediatr* 1975; 87:582-585
- Armstrong-Esther CA, Lacey JH, Crisp AH, et al: An investigation of the immune response of patients suffering from anorexia nervosa. *Postgrad Med* 1978; 54:395-399
- Crisp AH: Anorexia nervosa—Feeding disorder, nervous malnutrition or weight phobia. *World Rev Nutr Diet* 1970; 12:452-504
- Bowers TK, Eckert E: Leukopenia in anorexia nervosa. *Arch Intern Med* 1978; 138:1520-1523
- Deneke SM, Fanburg BL: Normobaric oxygen toxicity of the lung. *N Engl J Med* 1980; 303:76-86
- Deneke SM, Gershoff SN, Fanburg BL: Potentiation of oxygen toxicity in rats by dietary protein or amino acid deficiency. *J Appl Physiol* 1983; 54:147-151
- Donley AJ, Kemple TJ: Spontaneous pneumomediastinum complicating anorexia nervosa. *Br Med J* 1978; 2:1604-1605
- Garfinkel PE, Moldofsky H, Garner DM: The heterogeneity of anorexia nervosa as a distinct subgroup. *Arch Gen Psychiatry* 1980; 37:1036-1040
- Miller A, Bader RA, Bader ME, et al: Mineral oil pneumonia. *Ann Intern Med* 1962; 57:627-634
- Kennedy JD, Costello P, Balikian JP, et al: Exogenous lipid pneumonia. *AJR* 1981; 136:1145-1149
- Weill H, Ferrans VJ, Gay RM, et al: Early lipid pneumonia. *Am J Med* 1964; 36:370-376
- Bishop G: Oil aspiration pneumonia and pneumolipidosis. *Ann Intern Med* 1940; 23:1327-1359
- Volk BW, Nathanson L, Losner S, et al: Incidence of lipid pneumonia in a survey of 389 chronically ill patients. *Am J Med* 1951; 10:316-324
- Genereux GP: Lipids in the lungs: Radiologic-pathologic correlation. *J Can Assoc Radiol* 1970; 21:2-15
- Ayvazian LF, Steward DS, Merkel CG, et al: Diffuse lipid pneumonitis successfully treated with prednisone. *Am J Med* 1967; 43:930-934

Coexistent Dermatomyositis and Autoimmune Thyroiditis

THOMAS E. GAMSKY, MD
MICHAEL K. CHAN, MD
Martinez, California

DERMATOMYOSITIS is a disease characterized by nonsuppurative inflammation of skeletal muscle and characteristic skin changes. While dermatomyositis-like syndromes due to hypothyroidism have been well described, the occurrence of both hypothyroidism due to autoimmune thyroiditis and dermatomyositis together is not well documented.

We describe the case of a patient with coexistent autoimmune thyroiditis and dermatomyositis, showing clearly that these diseases can coexist. He was treated with pharmacologic doses of both steroids and thyroid hormone and is in clinical remission after two years.

Report of a Case

The patient, a 50-year-old male rice farmer, presented with proximal muscle weakness, Raynaud's phenomenon, dysphagia, hoarseness, and loss of energy for six months. On physical examination he had profound muscle weakness in the limb girdle and neck flexors. The patient was unable to lift his head from the pillow, rise from a chair, or walk up stairs without assistance. No fasciculations were seen. The neurologic examination elicited no abnormalities. Examination of the neck showed no goiter. His skin was remarkable for pronounced periungual erythema, a heliotrope rash over the upper eyelids, and Gottron's papules on the knuckles.

Admission laboratory tests were remarkable for the fol-

(Gamsky TE, Chan MK: Coexistent dermatomyositis and autoimmune thyroiditis. *West J Med* 1988 Feb; 148:213-214)

From the University of California, Davis, School of Medicine, and the Veterans Administration Medical Center, Martinez, California.

Reprint requests to Thomas E. Gamsky, MD, Division of Environmental and Occupational Medicine, LEHR Building, UCD School of Medicine, Davis, CA 95616.

ABBREVIATIONS USED IN TEXT

CK = creatine kinase
 EMG = electromyogram/phic
 TSH = thyroid-stimulating hormone

lowing values: creatine kinase (CK) 3,172 units per liter (normal 35 to 232); serum aspartate aminotransferase (glutamic-oxaloacetic transaminase) 112 units per liter (normal 0 to 41); lactate dehydrogenase 535 units per liter (normal 60 to 200); aldolase 28 mU per ml (normal 0.5 to 3.1); thyroxine 2.3 µg per dl (normal 5 to 12); free thyroxine index 0.9 (normal 1.7 to 5.4); thyroid-stimulating hormone (TSH) 45.1 IU per ml (normal 0.45 to 6.2); antinuclear antibody positive at 1:1,280 in a nucleolar pattern; antithyroid microsomal antibody positive at 1:1,600 dilution (normal less than 1:100); and antithyroglobulin antibody positive at 1:160 (normal less than 1:10).

A thyroid scan using radioactive iodine showed diffusely low uptake at five hours of 4.2% (normal 5% to 15%) and 24 hours' uptake at 10.6% (normal 12% to 36%). The thyroid size was estimated to be 20 grams. An erythrocyte sedimentation rate, chest x-ray film, electrocardiogram, complete blood count, anti-Smith antibody, anti-double-stranded DNA antibody, complement levels of C3 and C4, and antiribonuclear protein were all negative or normal. The test for the polymyositis-1 antigen was not done. An electromyogram (EMG) was diffusely abnormal with fibrillations, insertional irritability, positive sharp waves, and short-duration action potentials consistent with inflammatory myopathy. A biopsy specimen of the quadriceps showed perifascicular atrophy with lymphatic infiltrations of perifascicular collagen, muscle fascicle necrosis of type 1 and type 2 fibers, and basophilia consistent with polymyositis. An occult carcinoma workup was negative.

Clinical Course

Therapy consisting of the administration of levothyroxine sodium (Synthroid), 0.15 mg per day, and prednisone, 60 mg per day, was started. The CK level came down to 224 units per liter at one month and the TSH was down to 4.1 IU per ml. The prednisone dosage was rapidly tapered to 20 mg per day and at three months the CK value had returned to normal.

Clinically, the patient had quick resolution of his dysphagia, but muscle strength improved only slowly. Attempts to wean the prednisone therapy completely have been unsuccessful, and he is currently maintained on a regimen of 7.5 mg per day. At a two-year follow-up visit, his CK level and thyroid function tests remain normal, and muscle strength is adequate for farming chores.

Discussion

Hypothyroidism has long been associated with muscle weakness. Several authors have described a hypothyroid state clinically mimicking polymyositis/dermatomyositis.¹⁻⁶ Differentiating these diseases can be quite difficult, often relying on clinical grounds because laboratory test results can be ambiguous. We are aware of only one case report associating dermatomyositis and autoimmune thyroiditis.⁷ Our patient presented with the clinical features of dermatomyositis and was found to have coexistent hypothyroidism due to autoimmune thyroiditis.

This case shows the difficulties of interpreting laboratory data to diagnose these diseases. Muscle enzyme activity can

be normal or elevated in both diseases.⁸⁻¹⁰ Similarly, EMG studies in hypothyroid patients with myopathy have had a wide variety of findings, from fibrillations and positive sharp waves to normal results,^{8,10-14} although recent authors tend to support the view that the EMG should be normal in patients with hypothyroidism.¹³ Muscle biopsy specimens from hypothyroid patients have shown such diverse features as lymphocytic infiltrates, basophilia, and focal necrosis,^{1,14,15} typically seen in patients with polymyositis, while others have had no pathologic findings.^{15(pp642-643)} One conceivable explanation of these ambiguities lies in the possibility that some of those hypothyroid patients with abnormal EMG and muscle biopsy findings may have both diseases coexisting.

Immunologic studies have shown that both of these diseases are autoimmune mediated. Suppressor T-cell dysfunction and lymphocyte-mediated cytotoxicity have been well described in cases of polymyositis,¹⁶ and there is evidence of suppressor T-cell dysfunction in autoimmune thyroiditis as well,¹⁷⁻¹⁹ suggesting a common pathogenesis. In addition, the incidence of histocompatibility antigens HLA-B8 and DR3 is increased in frequency in autoimmune thyroiditis²⁰ and adult dermatomyositis.²¹ In our patient and in that of White and Tesar,⁷ hypothyroidism due to autoimmune thyroiditis most likely preceded the onset of dermatomyositis.

In conclusion, it is important for clinicians to be more aware that dermatomyositis and hypothyroidism due to autoimmune thyroiditis can coexist and a thorough evaluation of both may be necessary in patients presenting with proximal muscle weakness. In addition, further studies of the immunologic aspects of both diseases may unravel common pathways of autoimmune-mediated cellular dysfunction.

REFERENCES

1. Fessel WT: Myopathy of hypothyroidism. *Ann Rheum Dis* 1968; 27:590-596
2. Newman AJ, Lee C: Hypothyroidism simulating dermatomyositis. *J Pediatr* 1980; 95:772-774
3. Hochberg MC, Koppes GM, Edwards CQ, et al: Hypothyroidism presenting as a polymyositis-like syndrome—Report of two cases. *Arthritis Rheum* 1976; 19:1363-1366
4. Cabili S, Kaplinsky N, Pines A, et al: Hypothyroidism masquerading as polymyositis. *Postgrad Med J* 1982; 58:545-547
5. Mace BE, Stafforth JS, Mallaya AK: Severe myxedema presenting with chondrocalcinosis and polymyositis. *J R Soc Med* 1980; 73:887-888
6. Schnipper EF, Malamet RL, Arlin ZA: Polymyositis-like syndrome resulting from hypothyroidism in a patient with nodular lymphoma. *Clin Bull* 1981; 11:34-35
7. White SW, Tesar JT: Dermatomyositis and dermatitis herpetiformis. *Arch Dermatol* 1982; 118:599-601
8. Bohan A, Peter JB, Bowman RL, et al: Computer-assisted analysis of 153 patients with polymyositis and dermatomyositis. *Medicine (Baltimore)* 1977; 56:255-286
9. Hanson V, Kornreich H: Systemic rheumatic disorders ('collagen disease') in childhood: Lupus erythematosus, anaphylactoid purpura, dermatomyositis and scleroderma—I. *Bull Rheum Dis* 1967; 17:435-441
10. Vignos PJ Jr, Goldwyn J: Evaluation of laboratory tests in diagnosis and management of polymyositis. *Am J Med Sci* 1972; 263:291-308
11. Maracci AA: Electromyogram in diagnosis of polymyositis. *Electromyography* 1965; 5:235-238
12. Lambert EH, Sayre GP, Eaton LM: Electrical activity in muscle in polymyositis. *Trans Am Neurol Assoc* 1954; 79:64-69
13. Lennon JA, Ritchie AE: *Clinical Electromyography*. Marshfield, Mass, Pitman Publishing, 1983, pp 164-167
14. Ramsey I: *Thyroid Disease and Muscle Dysfunction*. Chicago, Year Book Medical Publishers, 1974, pp 138-149
15. Kakulac BA, Adams RD: *Diseases of Muscle*. Philadelphia, Harper & Row, 1985
16. Engel AG, Arahata K: Monoclonal antibody analysis of mononuclear cells in myopathies—II: Phenotypes of autoinvasive cells in polymyositis and inclusion body myositis. *Ann Neurol* 1984; 16:209-215
17. Solomon DH, Beall GN, Terasaki PI, et al: Autoimmune thyroid diseases—Graves' and Hashimoto's (UCLA Conference). *Ann Intern Med* 1978; 88:379-391
18. Strakosch CR: Immunology of autoimmune thyroid diseases. *N Engl J Med* 1982; 307:1499-1507
19. Wall JR, Fang SL, Walters BA, et al: Multisystem immunologically mediated disease: T lymphocyte deficiency and thyroid immunologic disease—A report of four cases. *Aust NZ J Med* 1978; 8:290-294
20. Walfish PG, Farid NR: The immunogenetic basis of autoimmune thyroid disease. In Walfish PG, Wall JR, Volpe R (Eds): *Autoimmunity and the Thyroid*. Orlando, Fla, Academic Press, 1985, pp 9-36
21. Hirsch TT, Enlow RW, Bias WB, et al: HLA-D related (DR) antigens in various kinds of myositis. *Hum Immunol* 1981; 3:181-186